

**Aranesp<sup>®</sup>**  
**(darbepoetin alfa)**  
**For Injection**

**DESCRIPTION**

Aranesp<sup>®</sup> is an erythropoiesis stimulating protein, closely related to erythropoietin, that is produced in Chinese hamster ovary (CHO) cells by recombinant DNA technology. Aranesp<sup>®</sup> is a 165-amino acid protein that differs from recombinant human erythropoietin in containing 5 N-linked oligosaccharide chains, whereas recombinant human erythropoietin contains 3 chains<sup>1</sup>. The two additional N-glycosylation sites result from amino acid substitutions in the erythropoietin peptide backbone. The additional carbohydrate chains increase the approximate molecular weight of the glycoprotein from 30,000 to 37,000 daltons. Aranesp<sup>®</sup> is formulated as a sterile, colorless, preservative-free protein solution for intravenous (IV) or subcutaneous (SC) administration.

**Single-dose vials** are available containing 25, 40, 60, 100, 150, 200, 300, or 500 mcg of Aranesp<sup>®</sup>.

**Single-dose prefilled syringes** and prefilled SureClick<sup>™</sup> autoinjectors are available containing 25, 40, 60, 100, 150, 200, 300, or 500 mcg of Aranesp<sup>®</sup>. To reduce the risk of accidental needlesticks to users, each prefilled syringe is equipped with a needle guard that covers the needle during disposal.

Single-dose vials, prefilled syringes and autoinjectors are available in two formulations that contain excipients as follows:

**Polysorbate solution** Each 1 mL contains 0.05 mg polysorbate 80, and is formulated at pH  $6.2 \pm 0.2$  with 2.12 mg sodium phosphate monobasic monohydrate, 0.66 mg sodium phosphate dibasic anhydrous, and 8.18 mg sodium chloride in Water for Injection, USP (to 1 mL).

**Albumin solution** Each 1 mL contains 2.5 mg albumin (human), and is formulated at pH  $6.0 \pm 0.3$  with 2.23 mg sodium phosphate monobasic monohydrate, 0.53 mg sodium phosphate dibasic anhydrous, and 8.18 mg sodium chloride in Water for Injection, USP (to 1 mL).

**CLINICAL PHARMACOLOGY**

**Mechanism of Action**

Aranesp<sup>®</sup> stimulates erythropoiesis by the same mechanism as endogenous erythropoietin. A primary growth factor for erythroid development, erythropoietin is produced in the kidney and released into the bloodstream in response to hypoxia. In responding to hypoxia, erythropoietin interacts with progenitor stem cells to increase red blood cell (RBC) production. Production of endogenous erythropoietin is impaired in patients with chronic renal failure (CRF), and erythropoietin deficiency is the primary cause of their anemia. Increased hemoglobin levels are not generally observed until 2 to 6 weeks after initiating treatment with Aranesp<sup>®</sup> (see **DOSE AND ADMINISTRATION: Dose Adjustment**). In patients with cancer receiving concomitant chemotherapy, the etiology of anemia is multifactorial.

**Pharmacokinetics**

*Adult Patients*

The pharmacokinetics of Aranesp<sup>®</sup> were studied in patients with CRF and cancer patients receiving chemotherapy.

Following intravenous (IV) administration in CRF patients, Aranesp<sup>®</sup> serum concentration-time profiles were biphasic, with a distribution half-life of approximately 1.4 hours and a mean terminal half-life of 21 hours. The terminal half-life of Aranesp<sup>®</sup> was approximately 3-fold longer than that of Epoetin alfa when administered intravenously.

Following subcutaneous (SC) administration, absorption is slow and rate limiting. The observed half-life in CRF patients, which reflected the rate of absorption, was 49 hours (range: 27 to 89 hours). Peak concentrations occurred at 34 hours (range: 24 to 72 hours). The bioavailability of Aranesp<sup>®</sup> as measured in CRF patients after SC administration was 37% (range: 30% to 50%).

Following the first SC dose of 6.75 mcg/kg (equivalent to 500 mcg for a 74-kg patient) in patients with cancer, the mean terminal half-life was 74 hours (range: 24 to 144 hours). Peak concentrations were observed at 90 hours (range: 71 to 123 hours) after a dose of 2.25 mcg/kg, and 71 hours (range: 28 to 120 hours) after a dose of 6.75 mcg/kg. When administered on a once-every-3-week (Q3W) schedule, 48-hour post-dose Aranesp<sup>®</sup> levels after the fourth dose were similar to those after the first dose.

Over the dose range of 0.45 to 4.5 mcg/kg Aranesp<sup>®</sup> administered IV or SC on a once-weekly (QW) schedule and 4.5 to 15 mcg/kg administered SC on a Q3W schedule, systemic exposure was approximately proportional to dose. No evidence of accumulation was observed beyond an expected < 2-fold increase in blood levels when compared to the initial dose.

#### *Pediatric Patients*

Aranesp<sup>®</sup> pharmacokinetics were studied in 12 pediatric CRF patients (age 3-16 years) receiving or not receiving dialysis. Following a single IV or SC Aranesp<sup>®</sup> dose, C<sub>max</sub> and half-life were similar to those obtained in adult CRF patients. Following a single SC dose, the average bioavailability was 54% (range: 32% to 70%), which was higher than that obtained in adult CRF patients.

## **CLINICAL STUDIES**

Throughout this section of the package insert, the Aranesp<sup>®</sup> study numbers associated with the nephrology and cancer clinical programs are designated with the letters “N” and “C”, respectively.

### ***Chronic Renal Failure Patients***

The safety and effectiveness of Aranesp<sup>®</sup> have been assessed in a number of multicenter studies. Two studies evaluated the safety and efficacy of Aranesp<sup>®</sup> for the correction of anemia in adult patients with CRF, and three studies (2 in adults and 1 in pediatric patients) assessed the ability of Aranesp<sup>®</sup> to maintain hemoglobin concentrations in patients with CRF who had been receiving other recombinant erythropoietins.

### **De Novo Use of Aranesp<sup>®</sup>**

In two open-label studies, Aranesp<sup>®</sup> or Epoetin alfa was administered for the correction of anemia in CRF patients who had not been receiving prior treatment with exogenous erythropoietin. Study N1 evaluated CRF patients receiving dialysis; Study N2 evaluated patients not requiring dialysis (predialysis patients). In both studies, the starting dose of Aranesp<sup>®</sup> was 0.45 mcg/kg administered once weekly. The starting dose of Epoetin alfa was 50 U/kg 3 times weekly in Study N1 and 50 U/kg twice weekly in Study N2. When necessary, dosage adjustments were instituted to maintain hemoglobin in the study target range of 11 to 13 g/dL. (Note: The recommended hemoglobin target is lower than the target range of these studies. See **DOSAGE AND ADMINISTRATION: General** for recommended clinical hemoglobin target.) The primary efficacy endpoint was the proportion of patients who experienced at least a 1.0 g/dL increase in

hemoglobin concentration to a level of at least 11.0 g/dL by 20 weeks (Study N1) or 24 weeks (Study N2). The studies were designed to assess the safety and effectiveness of Aranesp<sup>®</sup> but not to support conclusions regarding comparisons between the two products.

In Study N1, the hemoglobin target was achieved by 72% (95% CI: 62%, 81%) of the 90 patients treated with Aranesp<sup>®</sup> and 84% (95% CI: 66%, 95%) of the 31 patients treated with Epoetin alfa. The mean increase in hemoglobin over the initial 4 weeks of Aranesp<sup>®</sup> treatment was 1.10 g/dL (95% CI: 0.82 g/dL, 1.37 g/dL).

In Study N2, the primary efficacy endpoint was achieved by 93% (95% CI: 87%, 97%) of the 129 patients treated with Aranesp<sup>®</sup> and 92% (95% CI: 78%, 98%) of the 37 patients treated with Epoetin alfa. The mean increase in hemoglobin from baseline through the initial 4 weeks of Aranesp<sup>®</sup> treatment was 1.38 g/dL (95% CI: 1.21 g/dL, 1.55 g/dL).

### **Conversion From Other Recombinant Erythropoietins**

Two adult studies (N3 and N4) and one pediatric study (N5) were conducted in patients with CRF who had been receiving other recombinant erythropoietins. The studies compared the abilities of Aranesp<sup>®</sup> and other erythropoietins to maintain hemoglobin concentrations within a study target range of 9 to 13 g/dL in adults and 10 to 12.5 g/dL in pediatric patients. (Note: The recommended hemoglobin target is lower than the target range of these studies. See **DOSE AND ADMINISTRATION: General** for recommended clinical hemoglobin target.) CRF patients who had been receiving stable doses of other recombinant erythropoietins were randomized to Aranesp<sup>®</sup>, or to continue with their prior erythropoietin at the previous dose and schedule. For patients randomized to Aranesp<sup>®</sup>, the initial weekly dose was determined on the basis of the previous total weekly dose of recombinant erythropoietin.

#### *Adult Patients*

Study N3 was a double-blind study conducted in North America, in which 169 hemodialysis patients were randomized to treatment with Aranesp<sup>®</sup> and 338 patients continued on Epoetin alfa. Study N4 was an open-label study conducted in Europe and Australia in which 347 patients were randomized to treatment with Aranesp<sup>®</sup> and 175 patients were randomized to continue on Epoetin alfa or Epoetin beta. Of the 347 patients randomized to Aranesp<sup>®</sup>, 92% were receiving hemodialysis and 8% were receiving peritoneal dialysis.

In Study N3, a median weekly dose of 0.53 mcg/kg Aranesp<sup>®</sup> (25th, 75th percentiles: 0.30, 0.93 mcg/kg) was required to maintain hemoglobin in the study target range. In Study N4, a median weekly dose of 0.41 mcg/kg Aranesp<sup>®</sup> (25th, 75th percentiles: 0.26, 0.65 mcg/kg) was required to maintain hemoglobin in the study target range.

#### *Pediatric Patients*

Study N5 was an open-label, randomized study, conducted in the United States in pediatric patients from 1 to 18 years of age with CRF receiving or not receiving dialysis. Patients that were stable on Epoetin alfa were randomized to receive either darbepoetin alfa (n = 82) administered once weekly (SC or IV) or to continue receiving Epoetin alfa (n = 42) at the current dose, schedule, and route of administration. A median weekly dose of 0.41 mcg/kg Aranesp<sup>®</sup> (25th, 75th percentiles: 0.25, 0.82 mcg/kg) was required to maintain hemoglobin in the study target range.

### **Cancer Patients Receiving Chemotherapy**

#### *Once-Weekly (QW) Dosing*

The safety and effectiveness of Aranesp<sup>®</sup> in reducing the requirement for RBC transfusions in patients undergoing chemotherapy was assessed in a randomized, placebo-controlled, double-blind, multinational study (C1). This study was conducted in anemic (Hgb  $\leq$  11 g/dL) patients with advanced, small cell or non-small cell lung cancer, who received a platinum-containing chemotherapy regimen. Patients were randomized to receive Aranesp<sup>®</sup> 2.25 mcg/kg (n = 156) or placebo (n = 158) administered as a single weekly SC injection for up to 12 weeks. The dose was escalated to 4.5 mcg/kg/week at week 6, in subjects with an inadequate response to treatment, defined as less than 1 g/dL hemoglobin increase. There were 67 patients in the Aranesp<sup>®</sup> arm who had their dose increased from 2.25 to 4.5 mcg/kg/week, at any time during the treatment period.

Efficacy was determined by a reduction in the proportion of patients who were transfused over the 12-week treatment period. A significantly lower proportion of patients in the Aranesp<sup>®</sup> arm, 26% (95% CI: 20%, 33%) required transfusion compared to 60% (95% CI: 52%, 68%) in the placebo arm (Kaplan-Meier estimate of proportion;  $p < 0.001$  by Cochran-Mantel-Haenszel test). Of the 67 patients who received a dose increase, 28% had a 2 g/dL increase in hemoglobin over baseline, generally occurring between weeks 8 to 13. Of the 89 patients who did not receive a dose increase, 69% had a 2 g/dL increase in hemoglobin over baseline, generally occurring between weeks 6 to 13.

#### *Once-Every-3-Week (Q3W) Dosing*

The safety and effectiveness of Q3W Aranesp<sup>®</sup> therapy in reducing the requirement for red blood cell (RBC) transfusions in patients undergoing chemotherapy was assessed in a randomized, double-blind, multinational study (C2). This study was conducted in anemic (Hgb  $<$  11 g/dL) patients with non-myeloid malignancies receiving multicycle chemotherapy. Patients were randomized to receive Aranesp<sup>®</sup> at 500 mcg Q3W (n = 353) or 2.25 mcg/kg (n = 352) administered weekly as a SC injection for up to 15 weeks. In both groups, the dose was reduced by 40% of the previous dose (e.g., for first dose reduction, to 300 mcg in the Q3W group and 1.35 mcg/kg in the QW group) if hemoglobin increased by more than 1 g/dL in a 14-day period. Study drug was withheld if hemoglobin exceeded 13 g/dL. In the Q3W group, 254 patients (72%) required dose reductions (median time to first reduction at 6 weeks). In the QW group, 263 patients (75%) required dose reductions (median time to first reduction at 5 weeks).

Efficacy was determined by a comparison of the Kaplan-Meier estimates of the proportion of patients who received at least one RBC transfusion between day 29 and the end of treatment. Three hundred thirty five patients in the Q3W group and 337 patients in the QW group remained on study through or beyond day 29 and were evaluated for efficacy. Twenty seven percent (95% CI: 22%, 32%) of patients in the Q3W group and 34% (95% CI: 29%, 39%) in the weekly group required a RBC transfusion. The observed difference in the transfusion rates (Q3W-QW) was -6.7% (95% CI: -13.8%, 0.4%).

#### **INDICATIONS AND USAGE**

Aranesp<sup>®</sup> is indicated for the treatment of anemia associated with chronic renal failure, including patients on dialysis and patients not on dialysis, and for the treatment of anemia in patients with non-myeloid malignancies where anemia is due to the effect of concomitantly administered chemotherapy.

#### **CONTRAINDICATIONS**

Aranesp<sup>®</sup> is contraindicated in patients with:

- uncontrolled hypertension
- known hypersensitivity to the active substance or any of the excipients

## WARNINGS

### Mortality, Cardiovascular Events, and Hemoglobin Levels

**Aranesp<sup>®</sup> and other erythropoietic therapies may increase the risk of cardiovascular events, including death. The higher risk of cardiovascular events may be associated with higher hemoglobin and/or higher rates of rise of hemoglobin. The hemoglobin level should be managed carefully to avoid exceeding a target level of 12 g/dL.**

In a clinical trial of Epoetin alfa (rHuEPO) treatment in hemodialysis patients with clinically evident cardiac disease, patients were randomized to a target hemoglobin of either  $14 \pm 1$  g/dL or  $10 \pm 1$  g/dL<sup>2</sup>. Higher mortality (35% vs 29%) was observed in the 634 patients randomized to a target hemoglobin of 14 g/dL than in the 631 patients assigned a target hemoglobin of 10 g/dL. The reason for the increased mortality observed in this study is unknown; however, the incidence of nonfatal myocardial infarction, vascular access thrombosis, and other thrombotic events was also higher in the group randomized to a target hemoglobin of 14 g/dL.

In another clinical trial of Epoetin alfa, 1432 predialysis patients were prospectively randomized in an open-label manner to determine if treatment to a target hemoglobin of 13.5 g/dL would improve mortality and cardiovascular outcomes, compared to treatment to a target hemoglobin of 11.3 g/dL. The study was terminated after interim analysis showed little chance to demonstrate reduction of the composite endpoint rate for the higher hemoglobin group compared to the lower hemoglobin group. Final results showed that 125 of 715 patients (18%) randomized to the higher hemoglobin group reached the composite endpoint of death, myocardial infarction, stroke, and hospitalization for congestive heart failure, compared to 97 of 717 patients (14%) in the lower hemoglobin group (HR 1.3, 95% CI 1.0-1.7, p=0.03). Each endpoint was evaluated independently. In the higher hemoglobin group, there were 52 deaths (7%), 18 myocardial infarctions (3%), 64 hospitalizations for congestive heart failure (9%), and 12 strokes (2%) as compared to the lower hemoglobin group with 36 deaths (5%), 20 myocardial infarctions (3%), 47 hospitalizations for congestive heart failure (7%), and 12 strokes (2%). The high patient drop out rate (38%) in this study should be considered when interpreting the results.

In patients treated with Aranesp<sup>®</sup> or other recombinant erythropoietins in Aranesp<sup>®</sup> clinical trials, increases in hemoglobin greater than approximately 1.0 g/dL during any 2-week period were associated with increased incidence of cardiac arrest, neurologic events (including seizures and stroke), exacerbations of hypertension, congestive heart failure, vascular thrombosis/ischemia/infarction, acute myocardial infarction, and fluid overload/edema. It is recommended that the dose of Aranesp<sup>®</sup> be decreased if the hemoglobin increase exceeds 1.0 g/dL in any 2-week period, because of the association of excessive rate of rise of hemoglobin with these events.

### Hypertension

Patients with uncontrolled hypertension should not be treated with Aranesp<sup>®</sup>; blood pressure should be controlled adequately before initiation of therapy. Blood pressure may rise during treatment of anemia with Aranesp<sup>®</sup> or Epoetin alfa. In Aranesp<sup>®</sup> clinical trials, approximately 40% of patients with CRF required initiation or intensification of antihypertensive therapy during the early phase of treatment when the hemoglobin was increasing. Hypertensive encephalopathy and seizures have been observed in patients with CRF treated with Aranesp<sup>®</sup> or Epoetin alfa.

Special care should be taken to closely monitor and control blood pressure in patients treated with Aranesp<sup>®</sup>. During Aranesp<sup>®</sup> therapy, patients should be advised of the importance of compliance with antihypertensive therapy and dietary restrictions. If blood pressure is difficult to control by pharmacologic or dietary measures, the dose of Aranesp<sup>®</sup> should be reduced or withheld (see **DOSAGE AND ADMINISTRATION: Dose Adjustment**). A clinically significant decrease in hemoglobin may not be observed for several weeks.

## Seizures

Seizures have occurred in patients with CRF participating in clinical trials of Aranesp<sup>®</sup> and Epoetin alfa. During the first several months of therapy, blood pressure and the presence of premonitory neurologic symptoms should be monitored closely. While the relationship between seizures and the rate of rise of hemoglobin is uncertain, it is recommended that the dose of Aranesp<sup>®</sup> be decreased if the hemoglobin increase exceeds 1.0 g/dL in any 2-week period.

## Thrombotic Events and Increased Mortality

An increased incidence of thrombotic events has been observed in patients treated with erythropoietic agents. In patients with cancer who received Aranesp<sup>®</sup>, pulmonary emboli, thrombophlebitis and thrombosis occurred more frequently than in placebo controls (see **ADVERSE REACTIONS: Cancer Patients Receiving Chemotherapy, Table 4**).

In a randomized controlled study with another erythropoietic product in 939 women with metastatic breast cancer receiving chemotherapy, patients received either weekly Epoetin alfa or placebo for up to a year. This study was designed to prevent anemia (maintain hemoglobin levels between 12 and 14 g/dL or hematocrit between 36% and 42%). Treatment with Epoetin alfa was associated with a higher rate of fatal thrombotic events (1.1% Epoetin alfa vs 0.2% placebo) in the first 4 months of the study. Based on Kaplan-Meier estimates, the proportion of subjects surviving at 12 months after randomization was lower in the Epoetin alfa group than in the placebo group (70% vs 76%),  $p = 0.012$ , log rank. However, due to insufficient monitoring and data collection, reliable comparisons cannot be made concerning the effect of Epoetin alfa on overall time to disease progression, progression-free survival, and overall survival. Until further information is available, the recommended target hemoglobin should not exceed 12 g/dL in men or women.

## Pure Red Cell Aplasia

Cases of pure red cell aplasia (PRCA) and of severe anemia, with or without other cytopenias, associated with neutralizing antibodies to erythropoietin have been reported in patients treated with Aranesp<sup>®</sup>. This has been reported predominantly in patients with CRF receiving Aranesp<sup>®</sup> by subcutaneous administration. Any patient who develops a sudden loss of response to Aranesp<sup>®</sup>, accompanied by severe anemia and low reticulocyte count, should be evaluated for the etiology of loss of effect, including the presence of neutralizing antibodies to erythropoietin (see **PRECAUTIONS: Lack or Loss of Response to Aranesp<sup>®</sup>**). If anti-erythropoietin antibody-associated anemia is suspected, withhold Aranesp<sup>®</sup> and other erythropoietic proteins. Contact Amgen (1-800-77AMGEN) to perform assays for binding and neutralizing antibodies. Aranesp<sup>®</sup> should be permanently discontinued in patients with antibody-mediated anemia. Patients should not be switched to other erythropoietic proteins as antibodies may cross-react (see **ADVERSE REACTIONS: Immunogenicity**).

## Albumin (Human)

Aranesp<sup>®</sup> is supplied in two formulations with different excipients, one containing polysorbate 80 and another containing albumin (human), a derivative of human blood (see **DESCRIPTION**). Based on effective donor screening and product manufacturing processes, Aranesp<sup>®</sup> formulated with albumin carries an extremely remote risk for transmission of viral diseases. A theoretical risk for transmission of Creutzfeldt-Jakob disease (CJD) also is considered extremely remote. No cases of transmission of viral diseases or CJD have ever been identified for albumin.

## **PRECAUTIONS**

### **General**

The safety and efficacy of Aranesp<sup>®</sup> therapy have not been established in patients with underlying hematologic diseases (e.g., hemolytic anemia, sickle cell anemia, thalassemia, porphyria).

Aranesp<sup>®</sup> should be used with caution in patients with epilepsy.

The needle cover of the prefilled syringe contains dry natural rubber (a derivative of latex), which may cause allergic reactions in individuals sensitive to latex.

### **Lack or Loss of Response to Aranesp<sup>®</sup>**

A lack of response or failure to maintain a hemoglobin response with Aranesp<sup>®</sup> doses within the recommended dosing range should prompt a search for causative factors. Deficiencies of folic acid, iron or vitamin B<sub>12</sub> should be excluded or corrected. Depending on the clinical setting, intercurrent infections, inflammatory or malignant processes, osteofibrosis cystica, occult blood loss, hemolysis, severe aluminum toxicity and bone marrow fibrosis may compromise an erythropoietic response. In the absence of another etiology, the patient should be evaluated for evidence of PRCA and sera should be tested for the presence of antibodies to erythropoietin (see **WARNINGS: Pure Red Cell Aplasia**).

### **Hematology**

Sufficient time should be allowed to determine a patient's responsiveness to a dosage of Aranesp<sup>®</sup> before adjusting the dose. Because of the time required for erythropoiesis and the RBC half-life, an interval of 2 to 6 weeks may occur between the time of a dose adjustment (initiation, increase, decrease, or discontinuation) and a significant change in hemoglobin.

In order to prevent the hemoglobin from exceeding the recommended target (12 g/dL) or rising too rapidly (greater than 1.0 g/dL in 2 weeks), the guidelines for dose and frequency of dose adjustments should be followed (see **WARNINGS** and **DOSAGE AND ADMINISTRATION: Dose Adjustment**).

### **Allergic Reactions**

There have been rare reports of potentially serious allergic reactions, including skin rash and urticaria, associated with Aranesp<sup>®</sup>. Symptoms have recurred with rechallenge, suggesting a causal relationship exists in some instances. If a serious allergic or anaphylactic reaction occurs, Aranesp<sup>®</sup> should be immediately and permanently discontinued and appropriate therapy should be administered.

### **Patients with CRF Not Requiring Dialysis**

Patients with CRF not yet requiring dialysis may require lower maintenance doses of Aranesp<sup>®</sup> than patients receiving dialysis. Though predialysis patients generally receive less frequent monitoring of blood pressure and laboratory parameters than dialysis patients, predialysis patients may be more responsive to the effects of Aranesp<sup>®</sup>, and require judicious monitoring of blood pressure and hemoglobin. Renal function and fluid and electrolyte balance should also be closely monitored.

### **Dialysis Management**

Therapy with Aranesp<sup>®</sup> results in an increase in RBCs and a decrease in plasma volume, which could reduce dialysis efficiency; patients who are marginally dialyzed may require adjustments in their dialysis prescription.

### **Tumor Growth Factor Potential**

Aranesp<sup>®</sup> is a growth factor that primarily stimulates RBC production. Erythropoietin receptors are also found on the surfaces of normal, non-hematopoietic tissues and some malignant cell lines and tumor biopsy specimens. However, it is not known if these receptors are functional. The possibility that Aranesp<sup>®</sup> can act as a growth factor for any tumor type, particularly myeloid malignancies, has not been evaluated.

In a randomized, placebo-controlled study in 314 anemic subjects with advanced lung cancer randomized to either Aranesp<sup>®</sup> or placebo, statistically significant differences in time-to-progression (TTP) or overall survival (OS) were not observed; however, the study was not designed to detect or exclude clinically meaningful differences in either TTP or OS (see **CLINICAL STUDIES**).

Two additional studies explored the effect on survival and/or disease progression following administrations of two other erythropoietic products (ie, Epoetin alfa and Epoetin beta) with higher hemoglobin targets. The first study was a randomized controlled study in 939 women with metastatic breast cancer receiving chemotherapy where patients received either weekly Epoetin alfa or placebo for up to a year. This study was designed to prevent anemia (maintain hemoglobin levels between 12 and 14 g/dL or hematocrit between 36% and 42%). Mortality at 12 months was significantly higher in the Epoetin alfa arm (see **WARNINGS: Thrombotic Events and Increased Mortality**). This difference was observed primarily in the first 4 months of the study with more deaths attributed to breast cancer progression in the Epoetin alfa group (6% Epoetin alfa vs 3% placebo). Due to insufficient monitoring and data collection, reliable comparisons cannot be made concerning the effect of Epoetin alfa on overall time to disease progression, progression-free survival, and overall survival. The second study was a randomized controlled study in 351 head and neck cancer patients where Epoetin beta or placebo was administered to achieve target hemoglobins of 14 and 15 g/dL for women and men, respectively. Locoregional progression-free survival was significantly shorter (median of 406 days Epoetin beta vs 745 days placebo,  $p = 0.04$ ) in patients receiving Epoetin beta.

There is insufficient information to establish whether use of Epoetin products, including Aranesp<sup>®</sup>, have an adverse effect on time to tumor progression or progression-free survival.

These studies permitted or required dosing to achieve a hemoglobin level greater than 12 g/dL. Until further information is available, the recommended target hemoglobin should not exceed 12 g/dL in men or women.

### **Laboratory Tests**

After initiation of Aranesp<sup>®</sup> therapy, the hemoglobin should be determined weekly until it has stabilized and the maintenance dose has been established (see **DOSAGE AND ADMINISTRATION**). After a dose adjustment, the hemoglobin should be determined weekly for at least 4 weeks, until it has been determined that the hemoglobin has stabilized in response to the dose change. The hemoglobin should then be monitored at regular intervals.

In order to ensure effective erythropoiesis, iron status should be evaluated for all patients before and during treatment, as the majority of patients will eventually require supplemental iron therapy. Supplemental iron therapy is recommended for all patients whose serum ferritin is below 100 mcg/L or whose serum transferrin saturation is below 20%.

### **Information for Patients**

Patients should be informed of the possible side effects of Aranesp<sup>®</sup> and be instructed to report them to the prescribing physician. Patients should be informed of the signs and symptoms of allergic drug reactions and be advised of appropriate actions. Patients should be counseled on the importance of compliance with their Aranesp<sup>®</sup> treatment, dietary and dialysis prescriptions,

and the importance of judicious monitoring of blood pressure and hemoglobin concentration should be stressed.

It is recommended that Aranesp<sup>®</sup> should be administered by a healthcare professional. In those rare cases where it is determined that a patient can safely and effectively administer Aranesp<sup>®</sup> at home, appropriate instruction on the proper use of Aranesp<sup>®</sup> should be provided for patients and their caregivers, including careful review of the accompanying "Information for Patients" insert. Patients and caregivers should also be cautioned against the reuse of needles, syringes, prefilled SureClick<sup>™</sup> autoinjectors or drug product, and be thoroughly instructed in their proper disposal. A puncture-resistant container for the disposal of used syringes, autoinjectors, and needles should be made available to the patient. Patients should be informed that the needle cover on the prefilled syringe contains dry natural rubber (a derivative of latex), which should not be handled by persons sensitive to latex.

### **Drug Interactions**

No formal drug interaction studies of Aranesp<sup>®</sup> have been performed.

### **Carcinogenesis, Mutagenesis, and Impairment of Fertility**

**Carcinogenicity:** The carcinogenic potential of Aranesp<sup>®</sup> has not been evaluated in long-term animal studies. Aranesp<sup>®</sup> did not alter the proliferative response of non-hematological cells in vitro or in vivo. In toxicity studies of approximately 6 months duration in rats and dogs, no tumorigenic or unexpected mitogenic responses were observed in any tissue type. Using a panel of human tissues, the in vitro tissue binding profile of Aranesp<sup>®</sup> was identical to Epoetin alfa. Neither molecule bound to human tissues other than those expressing the erythropoietin receptor.

**Mutagenicity:** Aranesp<sup>®</sup> was negative in the in vitro bacterial and CHO cell assays to detect mutagenicity and in the in vivo mouse micronucleus assay to detect clastogenicity.

**Impairment of Fertility:** When administered intravenously to male and female rats prior to and during mating, reproductive performance, fertility and sperm assessment parameters were not affected at any doses evaluated (up to 10 mcg/kg/dose, administered 3 times weekly). An increase in post implantation fetal loss was seen at doses equal to or greater than 0.5 mcg/kg/dose, administered 3 times weekly.

### **Pregnancy Category C**

When Aranesp<sup>®</sup> was administered intravenously to rats and rabbits during gestation, no evidence of a direct embryotoxic, fetotoxic or teratogenic outcome was observed at doses up to 20 mcg/kg/day. The only adverse effect observed was a slight reduction in fetal weight, which occurred at doses causing exaggerated pharmacological effects in the dams (1 mcg/kg/day and higher). No deleterious effects on uterine implantation were seen in either species. No significant placental transfer of Aranesp<sup>®</sup> was observed in rats. An increase in post implantation fetal loss was observed in studies assessing fertility (see **PRECAUTIONS: Carcinogenesis, Mutagenesis, and Impairment of Fertility: Impairment of Fertility**).

Intravenous injection of Aranesp<sup>®</sup> to female rats every other day from day 6 of gestation through day 23 of lactation at doses of 2.5 mcg/kg/dose and higher resulted in offspring (F1 generation) with decreased body weights, which correlated with a low incidence of deaths, as well as delayed eye opening and delayed preputial separation. No adverse effects were seen in the F2 offspring.

There are no adequate and well-controlled studies in pregnant women. Aranesp<sup>®</sup> should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

## **Nursing Mothers**

It is not known whether Aranesp<sup>®</sup> is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when Aranesp<sup>®</sup> is administered to a nursing woman.

## **Pediatric Use**

### *Pediatric CRF Patients*

A study of the conversion from Epoetin alfa to Aranesp<sup>®</sup> among pediatric CRF patients over 1 year of age showed similar safety and efficacy to the findings from adult conversion studies (see **CLINICAL PHARMACOLOGY** and **CLINICAL STUDIES**). Safety and efficacy in the initial treatment of anemic pediatric CRF patients or in the conversion from another erythropoietin to Aranesp<sup>®</sup> in pediatric CRF patients less than 1 year of age have not been established.

### *Pediatric Cancer Patients*

The safety and efficacy of Aranesp<sup>®</sup> in pediatric cancer patients have not been established.

## **Geriatric Use**

Of the 1598 CRF patients in clinical studies of Aranesp<sup>®</sup>, 42% were age 65 and over, while 15% were age 75 and over. Of the 873 cancer patients in clinical studies receiving Aranesp<sup>®</sup> and concomitant chemotherapy, 45% were age 65 and over, while 14% were age 75 and over. No overall differences in safety or efficacy were observed between older and younger patients.

## **ADVERSE REACTIONS**

### **General**

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of Aranesp<sup>®</sup> cannot be directly compared to rates in the clinical trials of other drugs and may not reflect the rates observed in practice.

### **Immunogenicity**

As with all therapeutic proteins, there is a potential for immunogenicity. Neutralizing antibodies to erythropoietin, in association with PRCA or severe anemia (with or without other cytopenias), have been reported in patients receiving Aranesp<sup>®</sup> (see **WARNINGS: Pure Red Cell Aplasia**) during post-marketing experience.

In clinical studies, the percentage of patients with antibodies to Aranesp<sup>®</sup> was examined using the BIAcore assay. Sera from 1501 CRF patients and 1159 cancer patients were tested. At baseline, prior to Aranesp<sup>®</sup> treatment, binding antibodies were detected in 59 (4%) of CRF patients and 36 (3%) of cancer patients. While receiving Aranesp<sup>®</sup> therapy (range 22-177 weeks), a follow-up sample was taken. One additional CRF patient and eight additional cancer patients developed antibodies capable of binding Aranesp<sup>®</sup>. None of the patients had antibodies capable of neutralizing the activity of Aranesp<sup>®</sup> or endogenous erythropoietin at baseline or at end of study. No clinical sequelae consistent with PRCA were associated with the presence of these antibodies.

The incidence of antibody formation is highly dependent on the sensitivity and specificity of the assay. Additionally, the observed incidence of antibody (including neutralizing antibody) positivity in an assay may be influenced by several factors including assay methodology, sample handling, timing of sample collection, concomitant medications, and underlying disease. For these reasons, comparison of the incidence of antibodies across products within this class (erythropoietic proteins) may be misleading.

## **Chronic Renal Failure Patients**

### *Adult Patients*

In all studies, the most frequently reported serious adverse reactions with Aranesp<sup>®</sup> were vascular access thrombosis, congestive heart failure, sepsis, and cardiac arrhythmia. The most commonly reported adverse reactions were infection, hypertension, hypotension, myalgia, headache, and diarrhea (see **WARNINGS: Mortality, Cardiovascular Events, and Hemoglobin Levels and Hypertension**). The most frequently reported adverse reactions resulting in clinical intervention (e.g., discontinuation of Aranesp<sup>®</sup>, adjustment in dosage, or the need for concomitant medication to treat an adverse reaction symptom) were hypotension, hypertension, fever, myalgia, nausea, and chest pain.

The data described below reflect exposure to Aranesp<sup>®</sup> in 1598 CRF patients, including 675 exposed for at least 6 months, of whom 185 were exposed for greater than 1 year. Aranesp<sup>®</sup> was evaluated in active-controlled (n = 823) and uncontrolled studies (n = 775).

The rates of adverse events and association with Aranesp<sup>®</sup> are best assessed in the results from studies in which Aranesp<sup>®</sup> was used to stimulate erythropoiesis in patients anemic at study baseline (n = 348), and, in particular, the subset of these patients in randomized controlled trials (n = 276). Because there were no substantive differences in the rates of adverse reactions between these subpopulations, or between these subpopulations and the entire population of patients treated with Aranesp<sup>®</sup>, data from all 1598 patients were pooled.

The population encompassed an age range from 18 to 91 years. Fifty-seven percent of the patients were male. The percentages of Caucasian, Black, Asian, and Hispanic patients were 83%, 11%, 3%, and 1%, respectively. The median weekly dose of Aranesp<sup>®</sup> was 0.45 mcg/kg (25th, 75th percentiles: 0.29, 0.66 mcg/kg).

Some of the adverse events reported are typically associated with CRF, or recognized complications of dialysis, and may not necessarily be attributable to Aranesp<sup>®</sup> therapy. No important differences in adverse event rates between treatment groups were observed in controlled studies in which patients received Aranesp<sup>®</sup> or other recombinant erythropoietins.

The data in Table 1 reflect those adverse events occurring in at least 5% of patients treated with Aranesp<sup>®</sup>.

**Table 1. Adverse Events Occurring in  $\geq$  5% of CRF Patients**

Event	Patients Treated With Aranesp <sup>®</sup> (n = 1598)
<b>APPLICATION SITE</b>	
Injection-site Pain	7%
<b>BODY AS A WHOLE</b>	
Peripheral Edema	11%
Fatigue	9%
Fever	9%
Death	7%
Chest Pain, Unspecified	6%
Fluid Overload	6%
Access Infection	6%
Influenza-like Symptoms	6%
Access Hemorrhage	6%
Asthenia	5%
<b>CARDIOVASCULAR</b>	
Hypertension	23%
Hypotension	22%
Cardiac Arrhythmias/Cardiac Arrest	10%
Angina Pectoris/Cardiac Chest Pain	8%
Thrombosis Vascular Access	8%
Congestive Heart Failure	6%
<b>CNS/PNS</b>	
Headache	16%
Dizziness	8%
<b>GASTROINTESTINAL</b>	
Diarrhea	16%
Vomiting	15%
Nausea	14%
Abdominal Pain	12%
Constipation	5%
<b>MUSCULO-SKELETAL</b>	
Myalgia	21%
Arthralgia	11%
Limb Pain	10%
Back Pain	8%

(Continued)

**Table 1. Adverse Events Occurring in ≥ 5% of CRF Patients (Continued)**

Event	Patients Treated With Aranesp <sup>®</sup> (n = 1598)
RESISTANCE MECHANISM	
Infection <sup>a</sup>	27%
RESPIRATORY	
Upper Respiratory Infection	14%
Dyspnea	12%
Cough	10%
Bronchitis	6%
SKIN AND APPENDAGES	
Pruritus	8%

<sup>a</sup> Infection includes sepsis, bacteremia, pneumonia, peritonitis, and abscess.

The incidence rates for other clinically significant events are shown in Table 2.

**Table 2. Percent Incidence of Other Clinically Significant Events in CRF Patients**

Event	Patients Treated With Aranesp <sup>®</sup> (n = 1598)
Acute Myocardial Infarction	2%
Seizure	1%
Stroke	1%
Transient Ischemic Attack	1%

#### *Pediatric Patients*

In Study N5, Aranesp<sup>®</sup> was administered to 81 pediatric CRF patients who had stable hemoglobin concentrations while previously receiving Epoetin alfa (see **CLINICAL STUDIES**). In this study, the most frequently reported serious adverse reactions with Aranesp<sup>®</sup> were fever and dialysis access infection. The most commonly reported adverse reactions were fever, headache, upper respiratory infection, hypertension, hypotension, injection site pain and cough. Aranesp<sup>®</sup> administration was discontinued because of injection site pain in two patients and moderate hypertension in a third patient.

Studies have not evaluated the effects of Aranesp<sup>®</sup> when administered to pediatric patients as the initial treatment for the anemia associated with CRF.

#### **Thrombotic Events**

Vascular access thrombosis in hemodialysis patients occurred in clinical trials at an annualized rate of 0.22 events per patient year of Aranesp<sup>®</sup> therapy. Rates of thrombotic events (e.g., vascular access thrombosis, venous thrombosis, and pulmonary emboli) with Aranesp<sup>®</sup> therapy were similar to those observed with other recombinant erythropoietins in these trials; the median duration of exposure was 12 weeks.

### **Cancer Patients Receiving Chemotherapy**

The incidence data described below reflect the exposure to Aranesp® in 873 cancer patients including patients exposed to Aranesp® QW (547, 63%), Q2W (128, 16%), and Q3W (198, 23%). Aranesp® was evaluated in seven studies that were active-controlled and/or placebo-controlled studies of up to 6 months duration. The Aranesp®-treated patient demographics were as follows: median age of 63 years (range of 20 to 91 years); 40% male; 88% Caucasian, 5% Hispanic, 4% Black, and 3% Asian. Over 90% of patients had locally advanced or metastatic cancer, with the remainder having early stage disease. Patients with solid tumors (e.g., lung, breast, colon, ovarian cancers), and lymphoproliferative malignancies (e.g., lymphoma, multiple myeloma) were enrolled in the clinical studies. All of the 873 Aranesp®-treated subjects also received concomitant cyclic chemotherapy.

The most frequently reported serious adverse events included death (10%), fever (4%), pneumonia (3%), dehydration (3%), vomiting (2%), and dyspnea (2%). The most commonly reported adverse events were fatigue, edema, nausea, vomiting, diarrhea, fever and dyspnea (see **Table 3**). Except for those events listed in Tables 3 and 4, the incidence of adverse events in clinical studies occurred at a similar rate compared with patients who received placebo and were generally consistent with the underlying disease and its treatment with chemotherapy. The most frequently reported reasons for discontinuation of Aranesp® were progressive disease, death, discontinuation of the chemotherapy, asthenia, dyspnea, pneumonia and gastrointestinal hemorrhage. No important differences in adverse event rates between treatment groups were observed in controlled studies in which patients received Aranesp® or other recombinant erythropoietins.

**Table 3. Adverse Events Occurring in ≥ 5% of Patients Receiving Chemotherapy**

Event	Aranesp® (n = 873)	Placebo (n = 221)
<b>BODY AS A WHOLE</b>		
Fatigue	33%	30%
Edema	21%	10%
Fever	19%	16%
<b>CNS/PNS</b>		
Dizziness	14%	8%
Headache	12%	9%
<b>GASTROINTESTINAL</b>		
Diarrhea	22%	12%
Constipation	18%	17%
<b>METABOLIC/NUTRITION</b>		
Dehydration	5%	3%
<b>MUSCULO-SKELETAL</b>		
Arthralgia	13%	6%
Myalgia	8%	5%
<b>SKIN AND APPENDAGES</b>		
Rash	7%	3%

**Table 4. Incidence of Other Clinically Significant Adverse Events in Patients Receiving Chemotherapy**

Event	All Aranesp <sup>®</sup> (n = 873)	Placebo (n = 221)
Hypertension	3.7%	3.2%
Seizures/Convulsions <sup>a</sup>	0.6%	0.5%
Thrombotic Events	6.2%	4.1%
Pulmonary Embolism	1.3%	0.0%
Thrombosis <sup>b</sup>	5.6%	4.1%

<sup>a</sup> Seizures/Convulsions include the preferred terms: Convulsions, Convulsions Grand Mal, and Convulsions Local.

<sup>b</sup> Thrombosis includes: Thrombophlebitis, Thrombophlebitis Deep, Thrombosis Venous, Thrombosis Venous Deep, Thromboembolism, and Thrombosis.

In a randomized controlled trial of Aranesp<sup>®</sup> 500 mcg Q3W (n = 353) and Aranesp<sup>®</sup> 2.25 mcg/kg QW (n = 352), the incidences of all adverse events and of serious adverse events were similar between the two arms.

### Thrombotic and Cardiovascular Events

Overall, the incidence of thrombotic events was 6.2% for Aranesp<sup>®</sup> and 4.1% for placebo. However, the following events were reported more frequently in Aranesp<sup>®</sup>-treated patients than in placebo controls: pulmonary embolism, thromboembolism, thrombosis, and thrombophlebitis (deep and/or superficial). In addition, edema of any type was more frequently reported in Aranesp<sup>®</sup>-treated patients (21%) than in patients who received placebo (10%).

### OVERDOSAGE

The maximum amount of Aranesp<sup>®</sup> that can be safely administered in single or multiple doses has not been determined. Doses over 3.0 mcg/kg/week for up to 28 weeks have been administered to CRF patients. Doses up to 8.0 mcg/kg every week and 15.0 mcg/kg every 3 weeks have been administered to cancer patients for up to 12-16 weeks. Excessive rise and rate of rise in hemoglobin concentration, however, have been associated with adverse events (see **WARNINGS** and **DOSAGE AND ADMINISTRATION: Dose Adjustment**). In the event of polycythemia, Aranesp<sup>®</sup> should be temporarily withheld (see **DOSAGE AND ADMINISTRATION: Dose Adjustment**). If clinically indicated, phlebotomy may be performed.

### DOSAGE AND ADMINISTRATION

#### General

**IMPORTANT:** Aranesp<sup>®</sup> dosing regimens are different for each of the indications described in this section of the package insert. Aranesp<sup>®</sup> should be administered under the supervision of a healthcare professional.

Aranesp<sup>®</sup> is supplied in vials or in prefilled syringes with UltraSafe<sup>®</sup> Needle Guards<sup>†</sup>. Following administration of Aranesp<sup>®</sup> from the prefilled syringe, the UltraSafe<sup>®</sup> Needle Guard should be activated to prevent accidental needle sticks.

Aranesp<sup>®</sup> is also supplied in prefilled SureClick<sup>™</sup> autoinjectors containing the same dosage strengths as the prefilled syringes. Because the autoinjectors are designed to deliver the full content, autoinjectors should only be used for patients who need the full dose. If the required dose is not available in an autoinjector, prefilled syringes or vials should be used to administer the required dose. Autoinjectors are for subcutaneous administration only.

### **Chronic Renal Failure Patients**

Aranesp<sup>®</sup> is administered either IV or SC as a single weekly injection. ***In patients on hemodialysis, the IV route is recommended.*** The dose should be started and slowly adjusted as described below based on hemoglobin levels. If a patient fails to respond or maintain a response, this should be evaluated (see **WARNINGS: Pure Red Cell Aplasia**, **PRECAUTIONS: Lack or Loss of Response to Aranesp<sup>®</sup>** and **PRECAUTIONS: Laboratory Tests**). When Aranesp<sup>®</sup> therapy is initiated or adjusted, the hemoglobin should be followed weekly until stabilized and monitored at least monthly thereafter.

For patients who respond to Aranesp<sup>®</sup> with a rapid increase in hemoglobin (e.g., more than 1.0 g/dL in any 2-week period), the dose of Aranesp<sup>®</sup> should be reduced (see **DOSAGE AND ADMINISTRATION: Dose Adjustment**) because of the association of excessive rate of rise of hemoglobin with adverse events (see **WARNINGS: Mortality, Cardiovascular Events, and Hemoglobin Levels**).

The dose should be adjusted for each patient to achieve and maintain a target hemoglobin level not to exceed 12 g/dL.

### **Starting Dose**

#### **Correction of Anemia**

The recommended starting dose of Aranesp<sup>®</sup> for the correction of anemia in adult CRF patients is 0.45 mcg/kg body weight, administered as a single IV or SC injection once weekly. Because of individual variability, doses should be titrated to not exceed a target hemoglobin concentration of 12 g/dL (see **DOSAGE AND ADMINISTRATION: Dose Adjustment**). For many patients, the appropriate maintenance dose will be lower than this starting dose. Predialysis patients, in particular, may require lower maintenance doses. Also, some patients have been treated successfully with a SC dose of Aranesp<sup>®</sup> administered once every 2 weeks.

The use of Aranesp<sup>®</sup> in pediatric CRF patients as the initial treatment to correct anemia has not been studied.

#### **Conversion From Epoetin alfa to Aranesp<sup>®</sup>**

The starting weekly dose of Aranesp<sup>®</sup> for adults and pediatric patients should be estimated on the basis of the weekly Epoetin alfa dose at the time of substitution (see **Table 5**). For pediatric patients receiving a weekly Epoetin alfa dose of < 1500 units/week, the available data are insufficient to determine an Aranesp<sup>®</sup> conversion dose. Because of individual variability, doses should be titrated to maintain the target hemoglobin. Due to the longer serum half-life, Aranesp<sup>®</sup> should be administered less frequently than Epoetin alfa. Aranesp<sup>®</sup> should be administered once a week if a patient was receiving Epoetin alfa 2 to 3 times weekly. Aranesp<sup>®</sup> should be administered once every 2 weeks if a patient was receiving Epoetin alfa once per week. The route of administration (IV or SC) should be maintained.

**Table 5. Estimated Aranesp<sup>®</sup> Starting Doses (mcg/week) for Patients**

**Based on Previous Epoetin alfa Dose (Units/week)**

Previous Weekly Epoetin alfa Dose (Units/week)	Weekly Aranesp <sup>®</sup> Dose (mcg/week)	
	Adult	Pediatric
< 1,500	6.25	See text*
1,500 to 2,499	6.25	6.25
2,500 to 4,999	12.5	10
5,000 to 10,999	25	20
11,000 to 17,999	40	40
18,000 to 33,999	60	60
34,000 to 89,999	100	100
≥ 90,000	200	200

\*For pediatric patients receiving a weekly Epoetin alfa dose of < 1,500 units/week, the available data are insufficient to determine an Aranesp<sup>®</sup> conversion dose.

**Dose Adjustment**

The dose should be adjusted for each patient to achieve and maintain a target hemoglobin not to exceed 12 g/dL.

Increases in dose should not be made more frequently than once a month. If the hemoglobin is increasing and approaching 12 g/dL, the dose should be reduced by approximately 25%. If the hemoglobin continues to increase, doses should be temporarily withheld until the hemoglobin begins to decrease, at which point therapy should be reinitiated at a dose approximately 25% below the previous dose. If the hemoglobin increases by more than 1.0 g/dL in a 2-week period, the dose should be decreased by approximately 25%.

If the increase in hemoglobin is less than 1.0 g/dL over 4 weeks and iron stores are adequate (see **PRECAUTIONS: Laboratory Tests**), the dose of Aranesp<sup>®</sup> may be increased by approximately 25% of the previous dose. Further increases may be made at 4-week intervals until the specified hemoglobin is obtained.

**Maintenance Dose**

Aranesp<sup>®</sup> dosage should be adjusted to maintain a target hemoglobin not to exceed 12 g/dL. If the hemoglobin exceeds 12 g/dL, the dose may be adjusted as described above. Doses must be individualized to ensure that hemoglobin is maintained at an appropriate level for each patient.

**Cancer Patients Receiving Chemotherapy**

For pediatric patients, see **PRECAUTIONS: Pediatric Use**.

The recommended starting dose for Aranesp<sup>®</sup> administered weekly is 2.25 mcg/kg as a SC injection.

The recommended starting dose for Aranesp<sup>®</sup> administered once-every-3-weeks (Q3W) is 500 mcg as a SC injection.

For both dosing schedules, the dose should be adjusted for each patient to maintain a target hemoglobin not to exceed 12 g/dL. If the hemoglobin exceeds 13 g/dL, doses should be temporarily withheld until the hemoglobin falls to 12 g/dL. At this point, therapy should be reinitiated at a dose 40% below the previous dose. If the rate of hemoglobin increase is more than 1.0 g/dL per 2-week period or when the hemoglobin exceeds 11 g/dL, the dose should be reduced by 40% of the previous dose.

For patients receiving weekly administration, if there is less than a 1.0 g/dL increase in hemoglobin after 6 weeks of therapy, the dose of Aranesp<sup>®</sup> should be increased up to 4.5 mcg/kg.

### **Preparation and Administration of Aranesp<sup>®</sup>**

Do not shake Aranesp<sup>®</sup> or leave vials, syringes or prefilled SureClick<sup>™</sup> autoinjectors exposed to bright light. After removing the vials, prefilled syringes or autoinjectors from the cartons, keep them covered to protect from room light until administration. Vigorous shaking or exposure to light may denature Aranesp<sup>®</sup>, causing it to become biologically inactive. Always store vials, prefilled syringes or autoinjectors of Aranesp<sup>®</sup> in their carton until use.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration. Do not use any vials, prefilled syringes or autoinjectors exhibiting particulate matter or discoloration.

Do not dilute Aranesp<sup>®</sup>.

Do not administer Aranesp<sup>®</sup> in conjunction with other drug solutions.

Aranesp<sup>®</sup> contains no preservatives. Discard any unused portion. **Do not pool unused portions from the vials or prefilled syringes. Do not use the vial, prefilled syringe or autoinjector more than one time.**

Following administration of Aranesp<sup>®</sup> from the prefilled syringe, activate the UltraSafe<sup>®</sup> Needle Guard. Place your hands behind the needle, grasp the guard with one hand, and slide the guard forward until the needle is completely covered and the guard clicks into place. NOTE: If an audible click is not heard, the needle guard may not be completely activated.

The prefilled SureClick<sup>™</sup> autoinjector is designed to deliver the full dose. The completion of the injection is signaled by an audible click. Removal of the autoinjector from the injection site automatically extends a needle cover.

The autoinjectors, the syringes used with vials, and the entire prefilled syringe with activated needle guard should be disposed of in a puncture-proof container.

See the accompanying "Information for Patients" leaflet for complete instructions on the preparation and administration of Aranesp<sup>®</sup> for patients, including injection site selection.

### **HOW SUPPLIED**

Aranesp<sup>®</sup> is available in single-dose vials in two solutions, an albumin solution and a polysorbate solution. The words "Albumin Free" appear on the polysorbate container labels and the package main panels as well as other panels as space permits. Aranesp<sup>®</sup> single-dose prefilled syringes and prefilled SureClick<sup>™</sup> autoinjectors are available in albumin and polysorbate solutions. Both prefilled syringes and autoinjectors are supplied with a 27-gauge, ½-inch needle.

To reduce the risk of accidental needle sticks to users, each prefilled syringe is equipped with an UltraSafe® Needle Guard that is manually activated to cover the needle during disposal. The needle cover of the prefilled syringe contains dry natural rubber (a derivative of latex). The autoinjector has a needle cover that automatically extends as the autoinjector is removed from the injection site after completion of the injection.

Aranesp® is available in the following packages:

**Single-dose Vial, Polysorbate Solution**

<b>1 Vial/Pack, 4 Packs/Case</b>	<b>4 Vials/Pack, 4 Packs/Case</b>	<b>4 Vials/Pack, 10 Packs/Case</b>
200 mcg/1 mL (NDC 55513-006-01)	200 mcg/1 mL (NDC 55513-006-04)	25 mcg/1 mL (NDC 55513-002-04)
300 mcg/1 mL (NDC 55513-110-01)	300 mcg/1 mL (NDC 55513-110-04)	40 mcg/1 mL (NDC 55513-003-04)
500 mcg/1 mL (NDC 55513-008-01)		60 mcg/1 mL (NDC 55513-004-04)
		100 mcg/1 mL (NDC 55513-005-04)
		150 mcg/0.75 mL (NDC 55513-053-04)

**Single-dose Vial, Albumin Solution**

<b>1 Vial/Pack, 4 Packs/Case</b>	<b>4 Vials/Pack, 4 Packs/Case</b>	<b>4 Vials/Pack, 10 Packs/Case</b>
200 mcg/1 mL (NDC 55513-014-01)	200 mcg/1 mL (NDC 55513-014-04)	25 mcg/1 mL (NDC 55513-010-04)
300 mcg/1 mL (NDC 55513-015-01)	300 mcg/1 mL (NDC 55513-015-04)	40 mcg/1 mL (NDC 55513-011-04)
500 mcg/1 mL (NDC 55513-016-01)		60 mcg/1 mL (NDC 55513-012-04)
		100 mcg/1 mL (NDC 55513-013-04)
		150 mcg/0.75 mL (NDC 55513-054-04)

**Single-dose Prefilled Syringe (SingleJect®) with a 27-gauge, ½-inch needle with an UltraSafe® Needle Guard, Polysorbate Solution**

<b>1 Syringe/Pack, 4 Packs/Case</b>	<b>4 Syringes/Pack, 4 Packs/Case</b>	<b>4 Syringes/Pack, 10 Packs/Case</b>
200 mcg/0.4 mL (NDC 55513-028-01)	200 mcg/0.4 mL (NDC 55513-028-04)	25 mcg/0.42 mL (NDC 55513-057-04)
300 mcg/0.6 mL (NDC 55513-111-01)	300 mcg/0.6 mL (NDC 55513-111-04)	40 mcg/0.4 mL (NDC 55513-021-04)
500 mcg/1 mL (NDC 55513-032-01)		60 mcg/0.3 mL (NDC 55513-023-04)
		100 mcg/0.5 mL (NDC 55513-025-04)
		150 mcg/0.3 mL (NDC 55513-027-04)

**Single-dose Prefilled Syringe (SingleJect®) with a 27-gauge, ½-inch needle with an UltraSafe® Needle Guard, Albumin Solution**

<b>1 Syringe/Pack, 4 Packs/Case</b>	<b>4 Syringes/Pack, 4 Packs/Case</b>	<b>4 Syringes/Pack, 10 Packs/Case</b>
200 mcg/0.4 mL (NDC 55513-044-01)	200 mcg/0.4 mL (NDC 55513-044-04)	25 mcg/0.42 mL (NDC 55513-058-04)
300 mcg/0.6 mL (NDC 55513-046-01)	300 mcg/0.6 mL (NDC 55513-046-04)	40 mcg/0.4 mL (NDC 55513-037-04)
500 mcg/1 mL (NDC 55513-048-01)		60 mcg/0.3 mL (NDC 55513-039-04)
		100 mcg/0.5 mL (NDC 55513-041-04)
		150 mcg/0.3 mL (NDC 55513-043-04)

**Single-dose prefilled SureClick™ Autoinjector with a 27-gauge, 1/2-inch needle, Polysorbate Solution**

**1 Autoinjector/Pack**

25 mcg/0.42 mL  
(NDC 55513-090-01)

40 mcg/0.4 mL  
(NDC 55513-091-01)

60 mcg/0.3 mL  
(NDC 55513-092-01)

100 mcg/0.5 mL  
(NDC 55513-093-01)

150 mcg/0.3 mL  
(NDC 55513-094-01)

200 mcg/0.4 mL  
(NDC 55513-095-01)

300 mcg/0.6 mL  
(NDC 55513-096-01)

500 mcg/1 mL  
(NDC 55513-097-01)

**Single-dose prefilled SureClick™ Autoinjector with a 27-gauge, 1/2-inch needle, Albumin Solution**

**1 Autoinjector/Pack**

25 mcg/0.42 mL  
(NDC 55513-080-01)

40 mcg/0.4 mL  
(NDC 55513-081-01)

60 mcg/0.3 mL  
(NDC 55513-082-01)

100 mcg/0.5 mL  
(NDC 55513-083-01)

150 mcg/0.3 mL  
(NDC 55513-084-01)

200 mcg/0.4 mL  
(NDC 55513-085-01)

300 mcg/0.6 mL  
(NDC 55513-086-01)

500 mcg/1 mL  
(NDC 55513-087-01)

### **Storage**

**Store at 2° to 8°C (36° to 46°F). Do not freeze or shake. Protect from light.**

### **REFERENCES**

1. Egrie JC, Browne JK. Development and characterization of novel erythropoiesis stimulating protein (NESP). *Br J Cancer*. 2001;84 (suppl 1):3-10.
2. Besarab A, Bolton WK, Browne JK, et al. The effects of normal as compared with low hematocrit values in patients with cardiac disease who are receiving hemodialysis and epoetin. *N Engl J Med*. 1998; 339:584-590.

### **Rx only**

This product, or its use, may be covered by one or more US Patents, including US Patent No. 5,618,698, in addition to others including patents pending.

The logo for Amgen, featuring the word "AMGEN" in a bold, black, sans-serif font with a registered trademark symbol (®) to the upper right of the letter "N".

### **Manufactured by:**

Amgen Manufacturing, Limited, a subsidiary of Amgen Inc.  
One Amgen Center Drive  
Thousand Oaks, CA 91320-1799

©2001-2006 Amgen Inc. All rights reserved.

\* UltraSafe® is a registered trademark of Safety Syringes, Inc.

Issue Date: 12/15/2006

3xxxxxx – v13